



Sickle Cell Disease and COVID-19: An Outline to Decrease Burden and Minimize Morbidity

**Medical and Research Advisory Committee
Sickle Cell Disease Association of America**

This document will be updated once weekly on Friday afternoons as data and evidence emerge.

As of March 17, 2020, there are 188,421 infected and 7,499 deaths from COVID-19 worldwide. While most of these patients are in the Hubei province of China, the numbers in the United States are climbing exponentially with 4,748 cases and 93 deaths. The major morbidity and mortality from COVID-19 is described by Guan et al. NEJM 2020. In a cohort of 1099 patients, the median age of the patients was 47 years; 41.9% of the patients were female. Of this cohort, 5.0% were admitted to the ICU, 2.3% underwent invasive mechanical ventilation, and 1.4% died. The most common symptoms were fever (43.8% on admission and 88.7% during hospitalization) and cough (67.8%). On admission, ground-glass opacity was the most common radiologic finding on chest computed tomography (CT) (56.4%). No radiographic or CT abnormality was found in 157 of 877 patients (17.9%) with non-severe disease and in 5 of 173 patients (2.9%) with severe disease. It is clear from the work of our colleagues in China and Italy, that the disease is particularly devastating to individuals with pre-existing conditions.

Sickle cell disease (SCD) affects 100,000 individuals in the United States and many more world-wide. A major cause of morbidity and mortality in these individuals is acute chest syndrome (ACS) which is the term used for a constellation of findings that includes chest pain, cough, fever,

hypoxia (low oxygen level) and lung infiltrates. Acute chest syndrome may be the result of sickling in the small blood vessels, pulmonary infarction/emboli or viral or bacterial pneumonia. The management of ACS is challenging and requires vigilance from the medical team. There is a significant concern that the overlap of lung disease from COVID-19, in the setting of sickle cell lungs already primed for and scarred from acute chest syndrome may result in significant complications and amplification of healthcare utilization. Moreover, individuals with sickle cell disease have high utilization of acute care services including emergency departments and hospitals and often present with fever, signs and symptoms of pneumonia or evolving ACS, as well as acute sickle cell pain requiring parenteral therapy. Thus, there may be specific diagnostic, treatment and logistical challenges in meeting the healthcare needs of this population in the midst of the COVID-19 pandemic.

Here, we provide guidelines for providers who may encounter patients suspected of being infected with COVID-19 and having sickle cell disease.

Specifically, for sickle cell disease patients:

• **Clinic Flow and Strategies to Meet Transfusion Demands**

- Over the next several weeks, we will be inundated with persons under investigation (PUI) for COVID-19. Make every effort to interview the “person of interest” by telephone, text monitoring system, or video conference. Temperature monitoring could be reported by phone or shown to a provider via video conferencing.
- If possible, reschedule all routine appointments for the next 4-6 weeks. Push your institution to have virtual appointments and electronic refills.
- Reach out to your state board to have a waiver on duration of opioid prescriptions. (for example, the New York State Board has given three-month waivers to sickle cell patients).

- Halt all new research enrollment, including gene therapy/bone marrow transplantation
- Use enhanced discretion when ordering transfusions in sickle and non-sickle cell patients in the hospital and require all transfusions to be approved by hematology.
- Medical leaders should encourage local communities and political leadership to support local blood drives as blood shortages are anticipated.
- In the setting of blood shortage, clinicians will need to prioritize transfusion according to clinical need. Highest priorities include patients with recent stroke, progressive or critical neurovascular disease and those with repeated acute chest syndrome despite Hydroxyurea and significant cardiac or respiratory co-morbidity. Consider transitioning to Hydroxyurea for patients eligible according to TWITCH criteria. (Ware et al Lancet 2016)
- For patients with fever and cough
 - If possible at your center – test ALL patients for COVID-19.
 - Follow standard operating procedures for sickle cell disease and fever.
 - Consider minimizing hospitalizations if COVID-19 negative and manageable with outpatient antibiotics.

COVID-19+ Treatment in Patients with Sickle Cell Disease

- Consider early aggressive simple or exchange transfusion for patients with SCD and COVID-19 with fever and cough. This should certainly be done in patients with worsening anemia, evidence of hypoxia and chest x-ray changes, but may be done in the absence of these indications. This recommendation is based on H1N1 and sickle cell disease experience (Inusa et al Blood 2011, Jacobs et al PBC 2011).

- General respiratory measures for infected patients are to avoid aerosol-based interventions.
 - Nebulizers should not be used in a non-negative pressure room, instead use metered-dose inhaler for Albuterol
 - No non-invasive ventilation or high flow oxygen, or bronchoscopy on the general floors; should only be done in negative pressure rooms
 - Non-invasive oxygen therapy should progress to intubation to limit aerosolization and infection risk.
- Prophylactic antibiotic management and use oseltamivir until influenza ruled out
- Steroids are contraindicated until we have more data on immune system responses.
- Patients should be assessed for pulmonary hypertension and right ventricle dysfunction. Providers may consider inhaled nitric oxide in the setting of these entities.
- For severe and progressive cases, consider prone ventilation, inhaled nitric oxide (preferred) or Epoprostenol therapy, paralysis, and extra-corporeal membrane oxygenation.
- At the discretion of providers, Ritonavir/Lopinovir, Tocilizumab or Hydroxychloroquine may be used.
- The data around non-steroidal anti-inflammatory drugs, angiotensin converting enzyme inhibitors and angiotensin II receptor blockers are not clear and decisions about these drugs should be made on a case-by-case basis.

The coming months will be unprecedented in the history of healthcare. These recommendations serve as a template that may be beneficial when navigating this uncharted territory.

Authors listed below. In **Bold** are the authors responsible for the weekly updates. Their e-mails are available for suggestions/questions/comments.

SCDAA Medical and Research Advisory Committee Members

Miguel R Abboud, MD

Professor of Pediatrics and Pediatric
Hematology-Oncology
Chairman
Department of Pediatrics and Adolescent
Medicine
American University of Beirut, Lebanon

Biree Andemariam, MD

**Chair, Medical and Research Advisory
Committee, Sickle Cell Disease Association
of America
Chief Medical Officer, Sickle Cell Disease
Association of America
Director, New England Sickle Cell Institute
Associate Professor of Medicine
University of Connecticut Health
Farmington, CT 06030
andemariam@uchc.edu**

Shawn Bediako, PhD

Associate Professor
Department of Psychology
University of Maryland Baltimore County
Baltimore, Maryland

Andrew Campbell, MD

Center for Cancer and Blood Disorders
Children's National Health System
Associate Professor of Pediatrics
George Washington University School of
Medicine and Health Sciences

Raffaella Colombatti, MD, PhD

**Physician Azienda Ospedaliera-Università
di Padova
Department of Womens' and Child Health
Clinic of Pediatric Hematology Oncology
Via Giustiniani 3
35129 Padova Italy
raffaella.colombatti@aopd.veneto.it**

Lori Crosby, PsyD

Co-Director, Innovations in Community
Research, Division of Behavioral Medicine &
Clinical Psychology
Co-Director, CCTST, Community
Engagement Core
Psychologist, Research, Behavioral Medicine
& Clinical Psychologist
Cincinnati Children's
Professor, UC Department of Pediatrics

Deepika Darbari, MD

Center for Cancer and Blood Disorders
Children's National Health System
Associate Professor of Pediatrics
George Washington University School of
Medicine and Health Sciences

Payal Desai, MD

Associate Professor
Director of Sickle Cell Research
The Ohio State University
JamesCare at Ohio State East Hospital
181 Taylor Avenue
Columbus, Ohio 43203

James Eckman, MD

Professor Emeritus, Hematology & Medical
Oncology
Emory University School of Medicine
Department of Hematology and Medical
Oncology
49 Jesse Hill Jr. Drive SE
Atlanta, GA 30303

Mark Gladwin, MD

Professor and Chair
Department of Medicine
Founder, Pittsburgh Heart, Lung, and Blood
Vascular Medicine Institute
University of Pittsburgh
E1240 BST
200 Lothrop Street
Pittsburgh, PA 15261

Jo Howard, MB Bchir, MRCP, FRCPath

Head of Red Cell/Sickle Cell Service
Guy's and St Thomas'
NHS Foundation Trust
Great Maze Pond
London, United Kingdom

Lewis Hsu, MD, PhD

**Co-Chair, Medical and Research Advisory
Committee, Sickle Cell Disease Association
of America**
**Vice Chief Medical Officer, Sickle Cell
Disease Association of America**
Director of Pediatric Sickle Cell
**Professor of Pediatric Hematology-
Oncology**
University of Illinois at Chicago
Chicago, Illinois
lewhsu@uic.edu

Baba Inusa, MRCP, FRCPCH, DCP (Haem)

Honorary Reader in Paediatric
Haematology, King's College London
Lead Consultant Sickle Cell and
Thalassaemia Service
Evelina London Children's Hospital
St Thomas' Hospital
Westminster Bridge Road
London, United Kingdom

Elizabeth S. Klings, MD

Associate Professor of Medicine
Director, Center for Excellence in Sickle Cell
Disease
Director, Pulmonary Hypertension Center
Boston University School of Medicine
72 East Concord Street, R-304
Boston, MA 02118

Lakshmanan Krishnamurti, MD

Professor of Pediatrics, Director of BMT
Joseph Kuechenmeister Aflac Field Force
Chair
Aflac Cancer and Blood Disorders Center
Children's Healthcare of Atlanta/Emory
University

Sophie Lanzkron, MD, MHS

Director, Sickle Cell Center for Adults
The Johns Hopkins Hospital
1800 Orleans St
Baltimore, MD 21287

Caterina Minniti, MD

Director, Sickle Cell Center
Montefiore Health System
Professor of Medicine and Pediatrics
Albert Einstein College of Medicine
cminniti@montefiore.org

Genice T. Nelson, DNP, APRN, ANP-BC

Program Director
New England Sickle Cell Institute &
Connecticut Bleeding Disorders Programs
UConn Health
Farmington, CT 06030
Board Member, Sickle Cell Disease
Association of America

**Isaac Odame, MB ChB, MRCP(UK), FRCPath,
FRCPC, FRCPC**

Professor, Department of Paediatrics
University of Toronto
The Hospital for Sick Children
Division of Haematology/Oncology
555 University Avenue
Toronto, Ontario M5G 1X8

Gwendolyn Poles, D.O.

Honorary Medical Staff Member
Former Medical Director, Kline Health
Center
Faculty, Internal Medicine Program
UPMC Pinnacle
Harrisburg, PA
Board Member, Sickle Cell Disease
Association of America

John Roberts, MD

Yale Adult Sickle Cell Program
Smilow Cancer Hospital at Yale New Haven
35 Park Street, Ste 7th floor, Multispecialty
New Haven, CT 06511

Wally Smith, MD

Professor
Scientific Director, VCU Center on Health
Disparities
Director, VCU Adult Sickle Cell Program
Department of Internal Medicine
Division of General Internal Medicine
West Hospital W10W-403
P.O. Box 980306
Richmond, VA 23298

Crawford J Strunk MD

Pediatric Hematology/Oncology
Pediatric Hematology/Oncology Program at
Toledo Children's Hospital
2142 N. Cove Blvd.
Toledo, Ohio 43606

Immacolata Tartaglione, MD PhD

Department of Woman, Child and General
and Specialist Surgery
Università degli Studi della Campania "Luigi
Vanvitelli"
Naples, Italy

Marsha Treadwell, PhD

Director, Northern California, Network of
Care for Sickle Cell Disease
Co-Principal Investigator and Regional
Director, Pacific Sickle Cell Regional
Collaborative
Director, Hematology Behavioral Services
Comprehensive Center for Sickle Cell
Disease, Main Hospital
747 52nd Street
Oakland, CA 94609

Winfred C. Wang, MD

Member, Department of Hematology
St. Jude Children's Research Hospital
262 Danny Thomas Place, MS 800
Memphis, TN 38105

Russell E. Ware, MD, PhD

Director, Division of Hematology
Institute Co-Director, Cancer and Blood
Diseases Institute
Director, Global Health Center
Marjory J. Johnson Chair of Hematology
Translational Research
Cincinnati Children's
Professor, UC Department of Pediatrics

Julie Kanter Washko, MD

Associate Professor
Division of Hematology Oncology
University of Alabama at Birmingham
1720 2nd Avenue South, NP 2540
Birmingham, AL 35294-3300

Kim Smith-Whitley, MD

Professor of Pediatrics
Director Comprehensive Sickle Cell Center
Division of Hematology
11th Floor Colket Building
The Children's Hospital of Philadelphia
34th & Civic Center Blvd.
Philadelphia, PA 19104
Board Member, Sickle Cell Disease
Association of America

Wanda Whitten-Shurney, MD

CEO & Medical Director
Sickle Cell Disease Association, Michigan
Chapter Inc.
Board Member, Sickle Cell Disease
Association of America

Ahmar U. Zaidi, MD

**Assistant Professor of Pediatrics
Comprehensive Sickle Cell Center
Children's Hospital of Michigan
Director of Physician Network
Development, University Pediatricians
Wayne State University/Central Michigan
University School of Medicine
Detroit, MI
Ahmar@wayne.edu**